

STAT 3202: Practice 06

Autumn 2018, OSU

Exercise 1

Consider a random sample of size $n = 50$. The sample mean and standard deviation are:

- $\bar{x} = 5$
- $s = 3$

Use this sample to test $H_0: \mu = 4$ vs $H_1: \mu > 4$.

Report:

- The **test statistic**
 - The **critical value** when $\alpha = 0.05$.
 - A **decision** when $\alpha = 0.05$.
-

Exercise 2

Consider a random sample of size $n = 100$. The sample mean and standard deviation are:

- $\bar{x} = -0.5$
- $s = 2$

Use this sample to test $H_0: \mu = 0$ vs $H_1: \mu \neq 0$.

Report:

- The **test statistic**
 - The **p-value**
 - A **decision** when $\alpha = 0.01$.
-

Exercise 3

Consider two independent random samples.

Sample 1, from Population X :

- $n_x = 50$
- $\bar{x} = 5$
- $s_x = 2$

Sample 2, from Population Y :

- $n_y = 45$
- $\bar{y} = 6$
- $s_y = 3$

Use these samples to test $H_0: \mu_x = \mu_y$ vs $H_1: \mu_x \neq \mu_y$.

Report:

- The **test statistic**
 - The **p-value**
 - A **decision** when $\alpha = 0.05$.
-

Exercise 4

Consider two independent random samples.

Sample 1, from Population X :

- $n_x = 75$
- $\bar{x} = 13$
- $s_x = 5$

Sample 2, from Population Y :

- $n_y = 100$
- $\bar{y} = 11$
- $s_y = 6$

Use these samples to test $H_0: \mu_x = \mu_y$ vs $H_1: \mu_x \neq \mu_y$.

Report:

- The **test statistic**
 - The **critical values** when $\alpha = 0.05$.
 - A **decision** when $\alpha = 0.05$.
-

Exercise 5

Consider a random sample of size $n = 50$ from a dichotomous population. The sample proportion of the “success” class is

- $\hat{p} = 0.58$

Use this sample to test $H_0: p = 0.50$ vs $H_1: p \neq 0.50$.

Report:

- The **test statistic**
 - The **critical value** when $\alpha = 0.01$.
 - A **decision** when $\alpha = 0.01$.
-

Exercise 6

Consider a random sample of size $n = 100$ from a dichotomous population. The sample proportion of the “success” class is

- $\hat{p} = 0.81$

Use this sample to test $H_0: p = 0.70$ vs $H_1: p > 0.70$.

Report:

- The **test statistic**
 - The **p-value**
 - A **decision** when $\alpha = 0.05$.
-

Exercise 7

Consider two independent random samples from dichotomous populations.

Sample 1, from Population X :

- $n_x = 80$
- $\hat{p}_x = 0.70$

Sample 2, from Population Y :

- $n_y = 90$
- $\hat{p}_y = 0.79$

Use these samples to test $H_0: p_x = p_y$ vs $H_1: p_x \neq p_y$.

Report:

- The **test statistic**
 - The **p-value**
 - A **decision** when $\alpha = 0.10$.
-

Exercise 8

Consider two independent random samples from dichotomous populations.

Sample 1, from Population X :

- $n_x = 100$
- $\hat{p}_x = 0.39$

Sample 2, from Population Y :

- $n_y = 200$
- $\hat{p}_y = 0.51$

Use these samples to test $H_0: p_x = p_y$ vs $H_1: p_x \neq p_y$.

Report:

- The **test statistic**
 - The **p-value**
 - A **decision** when $\alpha = 0.01$.
-

Exercise 9

Consider a random sample of size $n = 12$ from a population that is assumed to be normal. The sample mean and standard deviation are:

- $\bar{x} = 5$
- $s = 2$

Use this sample to test $H_0: \mu = 4$ vs $H_1: \mu > 4$.

Report:

- The **test statistic**
 - The **critical value** when $\alpha = 0.05$.
 - A **decision** when $\alpha = 0.05$.
-

Exercise 10

Consider a random sample of size $n = 8$ from a population that is assumed to be normal. The sample mean and standard deviation are:

- $\bar{x} = -1.2$
- $s = 3$

Use this sample to test $H_0: \mu = 0$ vs $H_1: \mu < 0$.

Report:

- The **test statistic**
 - The **critical value** when $\alpha = 0.10$.
 - A **decision** when $\alpha = 0.10$.
-

Exercise 11

Consider a random sample of size $n = 12$ from a population that is assumed to be normal. The sample mean and standard deviation are:

- $\bar{x} = 5$
- $s = 2.3$

Use this sample to test $H_0: \sigma = 2$ vs $H_1: \sigma > 2$.

Report:

- The **test statistic**
 - The **critical value** when $\alpha = 0.05$.
 - A **decision** when $\alpha = 0.05$.
-

Exercise 12

Consider a random sample of size $n = 22$ from a population that is assumed to be normal. The sample mean and standard deviation are:

- $\bar{x} = 7$
- $s = 5.6$

Use this sample to test $H_0: \sigma = 5$ vs $H_1: \sigma > 5$.

Report:

- The **test statistic**
 - The **critical value** when $\alpha = 0.01$.
 - A **decision** when $\alpha = 0.01$.
-

Exercise 13

Consider two independent random samples. Assume both populations are normal and that their variances are equal.

Sample 1, from Population X :

- $n_x = 10$
- $\bar{x} = 5$
- $s_x = 2$

Sample 2, from Population Y :

- $n_y = 12$
- $\bar{y} = 6$
- $s_y = 1.5$

Use these samples to test $H_0: \mu_x = \mu_y$ vs $H_1: \mu_x \neq \mu_y$.

Report:

- The **test statistic**
 - The **critical values** when $\alpha = 0.05$.
 - A **decision** when $\alpha = 0.05$.
-

Exercise 14

Consider two independent random samples. Assume both populations are normal and that their variances are equal.

Sample 1, from Population X :

- $n_x = 14$
- $\bar{x} = 48,530$
- $s_x = 780$

Sample 2, from Population Y :

- $n_y = 11$

- $\bar{y} = 47,620$
- $s_y = 750$

Use these samples to test $H_0: \mu_x = \mu_y$ vs $H_1: \mu_x \neq \mu_y$.

Report:

- The **test statistic**
 - The **critical values** when $\alpha = 0.01$.
 - A **decision** when $\alpha = 0.01$.
-