

1. **2.3-11** **2.5-3** ~~2.5-3~~

If the moment-generating function of X is

$$M_X(t) = \frac{2}{5}e^t + \frac{1}{5}e^{2t} + \frac{2}{5}e^{3t},$$

Find the mean, variance, and pmf of X .

2. Suppose a discrete random variable X has the following probability distribution:

$$f(0) = \frac{7}{8}, \quad f(k) = \frac{1}{3^k}, \quad k = 2, 4, 6, 8, \dots$$

(possible values of X are even non-negative integers: $0, 2, 4, 6, 8, \dots$).

Recall Week 02 Discussion Problem 1 (a): this is a valid probability distribution.

- a) Find the moment-generating function of X , $M_X(t)$. For which values of t does it exist?
- b) Use $M_X(t)$ to find $E(X)$.

3. Suppose a discrete random variable X has the following probability distribution:

$$f(1) = \ln 3 - 1, \quad f(k) = \frac{(\ln 3)^k}{k!}, \quad k = 2, 3, 4, \dots$$

(possible values of X are positive integers: $1, 2, 3, 4, \dots$).

Recall Week 02 Discussion Problem 1 (b): this is a valid probability distribution.

- a) Find the moment-generating function of X , $M_X(t)$. For which values of t does it exist?
- b) Use $M_X(t)$ to find $E(X)$.

4. Suppose the moment-generating function of X is

$$M_X(t) = 0.1 e^{2t} + 0.3 e^{4t} + 0.6 e^{7t}.$$

- a) Find $\mu = E(X)$. b) Find $\sigma = SD(X)$.

5. Suppose a discrete random variable X has the following probability distribution:

$$f(k) = P(X = k) = a^k, \quad k = 2, 3, 4, 5, 6, \dots, \quad \text{zero otherwise.}$$

- a) Find the value of a that makes this is a valid probability distribution.
- b) Find $P(X \text{ is even})$.
- c) Find the moment-generating function of X , $M_X(t)$. For which values of t does it exist?
- d) Find $E(X)$.

6. Let X be a continuous random variable with the probability density function

$$f(x) = \frac{C}{x^4}, \quad x > 5, \quad \text{zero otherwise.}$$

- a) Find the value of C that would make $f(x)$ a valid probability density function.
- b) Find the cumulative distribution function of X , $F(x) = P(X \leq x)$.
 “Hint”: Should be $F(5) = 0$, $F(\infty) = 1$.
- c) Find the probability $P(6 < X < 10)$.
- f) Find the 80th percentile of the distribution of X , $\pi_{0.80}$.
- g) Find the expected value of X , $E(X)$.
- h) Find the standard deviation of X , $SD(X)$.

7. Let X be a continuous random variable with the probability density function

$$f(x) = Cx^2, \quad 3 \leq x \leq 9, \quad \text{zero otherwise.}$$

- a) Find the value of C that would make $f(x)$ a valid probability density function.
- b) Find the probability $P(X < 5)$.
- c) Find the probability $P(X > 7)$.
- d) Find the mean of the probability distribution of X .
- e) Find the median of the probability distribution of X .

8. Suppose a random variable X has the following probability density function:

$$f(x) = \cos x, \quad 0 < x < \frac{\pi}{2}, \quad \text{zero otherwise.}$$

- a) Find $P(X < \frac{\pi}{4})$.
- b) Find $\mu = E(X)$.
- c) Find the median of the probability distribution of X .

9. Let X be a continuous random variable with the probability density function

$$f(x) = 6x(1-x), \quad 0 < x < 1, \quad \text{zero elsewhere.}$$

Compute $P(\mu - 2\sigma < X < \mu + 2\sigma)$.

10. Suppose a random variable X has the following probability density function:

$$f(x) = xe^x, \quad 0 < x < 1, \quad \text{zero otherwise.}$$

- a) Find $P(X < \frac{1}{2})$.
- b) Find $\mu = E(X)$.
- c) Find the moment-generating function of X , $M_X(t)$.

11. Let X be a continuous random variable with the probability density function

$$f(x) = \begin{cases} c|x-3|, & 0 < x < 8, \\ 0, & \text{otherwise.} \end{cases}$$

- a) Find the value of c that makes $f(x)$ a valid probability density function.
- b) Find the probability $P(X < 5)$.
- c) Find the median of the probability distribution of X .
- d) Find the mean of the probability distribution of X .
- e) Find the variance of the probability distribution of X .

Answers:

1. **2.3-11** **2.5-3** ~~2.5-3~~

If the moment-generating function of X is

$$M_X(t) = \frac{2}{5}e^t + \frac{1}{5}e^{2t} + \frac{2}{5}e^{3t},$$

Find the mean, variance, and pmf of X .

$$f(1) = \frac{2}{5}, \quad f(2) = \frac{1}{5}, \quad f(3) = \frac{2}{5}.$$

$$M_X'(t) = \frac{2}{5}e^t + \frac{2}{5}e^{2t} + \frac{6}{5}e^{3t}. \quad E(X) = M_X'(0) = \mathbf{2}.$$

OR

$$E(X) = (1) \frac{2}{5} + (2) \frac{1}{5} + (3) \frac{2}{5} = \mathbf{2}.$$

$$M_X''(t) = \frac{2}{5}e^t + \frac{4}{5}e^{2t} + \frac{18}{5}e^{3t}. \quad E(X^2) = M_X''(0) = \frac{24}{5} = 4.8.$$

OR

$$E(X^2) = (1)^2 \frac{2}{5} + (2)^2 \frac{1}{5} + (3)^2 \frac{2}{5} = \frac{24}{5} = 4.8.$$

$$\text{Var}(X) = E(X^2) - [E(X)]^2 = 4.8 - 2^2 = \mathbf{0.8}.$$

2. Suppose a discrete random variable X has the following probability distribution:

$$f(0) = \frac{7}{8}, \quad f(k) = \frac{1}{3^k}, \quad k = 2, 4, 6, 8, \dots$$

(possible values of X are even non-negative integers: $0, 2, 4, 6, 8, \dots$).

Recall Discussion #2 Problem 1 (a): this is a valid probability distribution.

a) Find the moment-generating function of X , $M_X(t)$. For which values of t does it exist?

$$\begin{aligned} M_X(t) &= E(e^{tX}) = e^{0t} \cdot \frac{7}{8} + \sum_{k=1}^{\infty} e^{2kt} \cdot \frac{1}{3^{2k}} = \frac{7}{8} + \sum_{k=1}^{\infty} \left(\frac{e^{2t}}{9} \right)^k \\ &= \frac{7}{8} + \frac{\frac{e^{2t}}{9}}{1 - \frac{e^{2t}}{9}} = \frac{7}{8} + \frac{e^{2t}}{9 - e^{2t}} = \frac{9}{9 - e^{2t}} - \frac{1}{8}. \end{aligned}$$

$$\text{Must have } \left(\frac{e^{2t}}{9} \right) < 1 \text{ for geometric series to converge.} \quad \Rightarrow \quad t < \ln 3.$$

b) Use $M_X(t)$ to find $E(X)$.

$$\text{Recall Week 04 Discussion Problem 1 (a):} \quad E(X) = \frac{9}{32}.$$

$$M'_X(t) = \frac{2e^{2t}(9 - e^{2t}) - e^{2t}(-2e^{2t})}{(9 - e^{2t})^2} = \frac{18e^{2t}}{(9 - e^{2t})^2}, \quad t < \ln 3.$$

$$E(X) = M'_X(0) = \frac{18}{64} = \frac{9}{32}.$$

3. Suppose a discrete random variable X has the following probability distribution:

$$f(1) = \ln 3 - 1, \quad f(k) = \frac{(\ln 3)^k}{k!}, \quad k = 2, 3, 4, \dots$$

(possible values of X are positive integers: 1, 2, 3, 4, ...).

Recall Discussion #2 Problem 1 (b): this is a valid probability distribution.

“Hint”: Recall that $\sum_{k=0}^{\infty} \frac{a^k}{k!} = e^a$.

a) Find the moment-generating function of X , $M_X(t)$. For which values of t does it exist?

$$\begin{aligned} M_X(t) &= \sum_{\text{all } x} e^{tx} \cdot p(x) = e^{t \cdot (\ln 3 - 1)} + \sum_{k=2}^{\infty} e^{tk} \cdot \frac{(\ln 3)^k}{k!} \\ &= e^{t \cdot (\ln 3 - 1)} + \sum_{k=2}^{\infty} \frac{(e^t \ln 3)^k}{k!} = e^t \ln 3 - e^t + e^{e^t \ln 3} - 1 - e^t \ln 3 \\ &= 3e^t - e^t - 1, \quad t \in \mathbf{R}. \end{aligned}$$

b) Use $M_X(t)$ to find $E(X)$.

Recall Week 04 Discussion Problem 1 (b): $E(X) = 3 \ln 3 - 1 \approx 2.2958$.

$$M_X'(t) = 3e^t \cdot \ln 3 \cdot e^t - e^t, \quad E(X) = M_X'(0) = \mathbf{3 \ln 3 - 1}.$$

4. Suppose the moment-generating function of X is

$$M_X(t) = 0.1 e^{2t} + 0.3 e^{4t} + 0.6 e^{7t}.$$

a) Find $\mu = E(X)$.

$$M'_X(t) = 0.2 e^{2t} + 1.2 e^{4t} + 4.2 e^{7t}.$$

$$E(X) = M'_X(0) = 0.2 + 1.2 + 4.2 = \mathbf{5.6}.$$

b) Find $\sigma = SD(X)$.

$$M''_X(t) = 0.4 e^{2t} + 4.8 e^{4t} + 29.4 e^{7t}.$$

$$E(X^2) = M''_X(0) = 0.4 + 4.8 + 29.4 = 34.6.$$

$$\text{Var}(X) = E(X^2) - [E(X)]^2 = 34.6 - 5.6^2 = 3.24.$$

$$SD(X) = \sqrt{3.24} = \mathbf{1.8}.$$

OR

$$M_X(t) = 0.1 e^{2t} + 0.3 e^{4t} + 0.6 e^{7t}. \quad \Rightarrow$$

x	$f(x)$
2	0.1
4	0.3
7	0.6

x	$f(x)$	$x \cdot f(x)$	$x^2 \cdot f(x)$	$(x - \mu)^2 \cdot f(x)$
2	0.1	0.2	0.4	1.296
4	0.3	1.2	4.8	0.768
7	0.6	4.2	29.4	1.176
		5.6	34.6	3.240

$$\mu = E(X) = \sum x \cdot f(x) = \mathbf{5.6}. \quad \text{Var}(X) = \sum (x - \mu)^2 \cdot f(x) = 3.24.$$

$$\text{OR} \quad \text{Var}(X) = \sum x^2 \cdot f(x) - \mu^2 = 34.6 - 5.6^2 = 3.24.$$

$$SD(X) = \sqrt{3.24} = \mathbf{1.8}.$$

5. Suppose a discrete random variable X has the following probability distribution:

$$f(k) = P(X = k) = a^k, \quad k = 2, 3, 4, 5, 6, \dots, \quad \text{zero otherwise.}$$

a) Find the value of a that makes this is a valid probability distribution.

Must have $\sum_{\text{all } x} f(x) = 1.$

$$\Rightarrow 1 = \sum_{k=2}^{\infty} a^k = \frac{\text{first term}}{1 - \text{base}} = \frac{a^2}{1 - a}.$$

$$\Rightarrow a^2 + a - 1 = 0. \quad \Rightarrow a = \frac{-1 \pm \sqrt{5}}{2}.$$

$$\frac{-1 - \sqrt{5}}{2} < 0. \quad \Rightarrow a = \frac{\sqrt{5} - 1}{2} \approx 0.618034.$$

Note: $a = \frac{1}{\varphi} = \varphi - 1,$ where φ is the golden ratio.

b) Find $P(X \text{ is even})$.

$$P(X \text{ is even}) = f(2) + f(4) + f(6) + f(8) + \dots$$

$$= a^2 + a^4 + a^6 + a^8 + \dots = \frac{\text{first term}}{1 - \text{base}}$$

$$= \frac{a^2}{1 - a^2} = \frac{a^2}{a} = a \approx 0.618034.$$

- c) Find the moment-generating function of X , $M_X(t)$. For which values of t does it exist?

$$M_X(t) = E(e^{tX}) = \sum_{k=2}^{\infty} e^{tk} \cdot a^k = \sum_{k=2}^{\infty} (a \cdot e^t)^k = \frac{\text{first term}}{1 - \text{base}} = \frac{a^2 \cdot e^{2t}}{1 - a \cdot e^t},$$

$$(a \cdot e^t) < 1 \quad \Leftrightarrow \quad t < \ln \frac{1}{a} = \ln \varphi \approx 0.48121.$$

- d) Find $E(X)$.

$$M'_X(t) = \frac{2a^2 \cdot e^{2t} (1 - a \cdot e^t) - a^2 \cdot e^{2t} (-a \cdot e^t)}{(1 - a \cdot e^t)^2} = \frac{2a^2 \cdot e^{2t} - a^3 \cdot e^{3t}}{(1 - a \cdot e^t)^2},$$

$$t < \ln \frac{1}{a}.$$

$$E(X) = M'_X(0) = \frac{2a^2 - a^3}{(1-a)^2} = \frac{2-a}{1-a} = 3 + a \approx 3.618034.$$

OR

$$E(X) = 2 \cdot a^2 + 3 \cdot a^3 + 4 \cdot a^4 + 5 \cdot a^5 + 6 \cdot a^6 + \dots$$

$$a \cdot E(X) = 2 \cdot a^3 + 3 \cdot a^4 + 4 \cdot a^5 + 5 \cdot a^6 + \dots$$

$$\Rightarrow (1-a) \cdot E(X) = a^2 + a^2 + a^3 + a^4 + a^5 + a^6 + \dots = a^2 + 1.$$

$$\text{Therefore, } E(X) = \frac{a^2 + 1}{1-a} = \frac{2-a}{1-a} = 1 + \frac{1}{1-a} = 3 + a \approx 3.618034.$$

6. Let X be a continuous random variable with the probability density function

$$f(x) = \frac{C}{x^4}, \quad x > 5, \quad \text{zero otherwise.}$$

- a) Find the value of C that would make $f(x)$ a valid probability density function.

Must have $\int_{-\infty}^{\infty} f(x) dx = 1.$

$$1 = \int_5^{\infty} \frac{C}{x^4} dx = -\frac{C}{3x^3} \Big|_5^{\infty} = \frac{C}{375}. \quad C = \mathbf{375}.$$

- b) Find the cumulative distribution function of X , $F(x) = P(X \leq x)$.

“Hint”: Should be $F(5) = 0$, $F(\infty) = 1$.

$$F(x) = P(X \leq x) = \int_5^x \frac{375}{u^4} du = -\frac{375}{3u^3} \Big|_5^x = 1 - \frac{125}{x^3}, \quad x \geq 5.$$

- c) Find the probability $P(6 < X < 10)$.

$$\begin{aligned} P(6 < X < 10) &= \int_6^{10} \frac{375}{x^4} dx = -\frac{375}{3x^3} \Big|_6^{10} = \frac{125}{6^3} - \frac{125}{10^3} \\ &\approx 0.5787 - 0.1250 = \mathbf{0.4537}. \end{aligned}$$

OR

$$\begin{aligned} P(6 < X < 10) &= F(10-) - F(6) = \left(1 - \frac{125}{10^3}\right) - \left(1 - \frac{125}{6^3}\right) \\ &\approx 0.8750 - 0.4213 = \mathbf{0.4537}. \end{aligned}$$

f) Find the 80th percentile of the distribution of X , $\pi_{0.80}$.

$$P(X \leq \pi_{0.80}) = F(\pi_{0.80}) = 1 - \frac{125}{(\pi_{0.80})^3} = 0.80.$$

$$\pi_{0.80} = \sqrt[3]{625} \approx \mathbf{8.55}.$$

OR

$$P(X \geq \pi_{0.80}) = 0.20.$$

$$\int_{\pi_{0.80}}^{\infty} \frac{375}{x^4} dx = -\frac{375}{3x^3} \Big|_{\pi_{0.80}}^{\infty} = \frac{125}{(\pi_{0.80})^3} = 0.20.$$

$$\pi_{0.80} = \sqrt[3]{625} \approx \mathbf{8.55}.$$

For fun:

$$\text{Median} = \sqrt[3]{250} \approx 6.3.$$

g) Find the expected value of X , $E(X)$.

$$E(X) = \int_{-\infty}^{\infty} x \cdot f(x) dx = \int_5^{\infty} x \cdot \frac{375}{x^4} dx = \int_5^{\infty} \frac{375}{x^3} dx = -\frac{375}{2x^2} \Big|_5^{\infty} = \mathbf{7.5}.$$

h) Find the standard deviation of X , $SD(X)$.

$$E(X) = \int_{-\infty}^{\infty} x^2 \cdot f(x) dx = \int_5^{\infty} x^2 \cdot \frac{375}{x^4} dx = \int_5^{\infty} \frac{375}{x^2} dx = -\frac{375}{x} \Big|_5^{\infty} = 75.$$

$$\text{Var}(X) = E(X^2) - [E(X)]^2 = 75 - 7.5^2 = 18.75.$$

$$SD(X) = \sqrt{18.75} \approx \mathbf{4.33}.$$

7. Let X be a continuous random variable with the probability density function

$$f(x) = Cx^2, \quad 3 \leq x \leq 9, \quad \text{zero otherwise.}$$

a) Find the value of C that would make $f(x)$ a valid probability density function.

$$1 = \int_3^9 Cx^2 dx = \left. \frac{Cx^3}{3} \right|_3^9 = \frac{729 - 27}{3} C = 234C. \quad \Rightarrow \quad C = \frac{1}{234}.$$

b) Find the probability $P(X < 5)$.

$$P(X < 5) = \int_3^5 \frac{x^2}{234} dx = \left. \frac{x^3}{702} \right|_3^5 = \frac{125 - 27}{702} = \frac{98}{702} = \frac{49}{351} \approx 0.1396.$$

c) Find the probability $P(X > 7)$.

$$P(X > 7) = \int_7^9 \frac{x^2}{234} dx = \left. \frac{x^3}{702} \right|_7^9 = \frac{729 - 343}{702} = \frac{386}{702} = \frac{193}{351} \approx 0.5499.$$

d) Find the mean of the probability distribution of X .

$$E(X) = \int_{-\infty}^{\infty} x \cdot f(x) dx = \int_3^9 x \cdot \frac{x^2}{234} dx = \frac{x^4}{936} \Big|_3^9 = \frac{6480}{936} = \frac{90}{13} \approx 6.923.$$

e) Find the median of the probability distribution of X .

$$F(x) = P(X \leq x) = \int_{-\infty}^x f(y) dy = \int_3^x \frac{y^2}{234} dy = \frac{y^3}{702} \Big|_3^x = \frac{x^3 - 27}{702},$$

$3 \leq x < 9.$

$$F(x) = P(X \leq x) = 0, \quad x < 3.$$

$$F(x) = P(X \leq x) = 1, \quad x \geq 9.$$

$$F(m) = \frac{1}{2}. \quad \frac{m^3 - 27}{702} = \frac{1}{2}.$$

$$m^3 = \frac{702}{2} + 27 = 378. \quad m = \sqrt[3]{378} \approx 7.23.$$

8. Suppose a random variable X has the following probability density function:

$$f(x) = \cos x, \quad 0 < x < \frac{\pi}{2}, \quad \text{zero otherwise.}$$

a) Find $P(X < \frac{\pi}{4})$.

$$P(X < \frac{\pi}{4}) = \int_0^{\pi/4} \cos x \, dx = (\sin x) \Big|_0^{\pi/4} = \sin \frac{\pi}{4} - 0 = \frac{\sqrt{2}}{2} \approx 0.7071.$$

b) Find $\mu = E(X)$.

$$\mu = E(X) = \int_0^{\pi/2} x \cdot \cos x \, dx = (x \cdot \sin x + \cos x) \Big|_0^{\pi/2} = \frac{\pi}{2} - 1 \approx 0.5708.$$

c) Find the median of the probability distribution of X .

$$F(x) = P(X \leq x) = \int_0^x \cos y \, dy = \sin x, \quad 0 < x < \frac{\pi}{2}.$$

$$\text{Median: } F(m) = P(X \leq m) = \frac{1}{2}.$$

$$\Rightarrow \sin m = \frac{1}{2}. \quad m = \frac{\pi}{6} \approx 0.5236.$$

9. Let X be a continuous random variable with the probability density function

$$f(x) = 6x(1-x), \quad 0 < x < 1, \quad \text{zero elsewhere.}$$

Compute $P(\mu - 2\sigma < X < \mu + 2\sigma)$.

$$\mu = E(X) = \int_0^1 x \cdot 6x(1-x) dx = \int_0^1 6x^2 dx - \int_0^1 6x^3 dx = \frac{6}{3} - \frac{6}{4} = \frac{1}{2} = 0.50.$$

$$E(X^2) = \int_0^1 x^2 \cdot 6x(1-x) dx = \int_0^1 6x^3 dx - \int_0^1 6x^4 dx = \frac{6}{4} - \frac{6}{5} = 0.30.$$

$$\sigma^2 = \text{Var}(X) = E(X^2) - [E(X)]^2 = 0.30 - 0.25 = 0.05.$$

$$\sigma = \sqrt{0.05} \approx 0.2236.$$

$$\mu - 2\sigma \approx 0.0528,$$

$$\mu + 2\sigma \approx 0.9472.$$

$$\begin{aligned} P(\mu - 2\sigma < X < \mu + 2\sigma) &\approx \int_{0.0528}^{0.9472} 6x(1-x) dx = \left(3x^2 - 2x^3 \right) \Big|_{0.0528}^{0.9472} \\ &\approx 0.992 - 0.008 = \mathbf{0.984}. \end{aligned}$$

10. Suppose a random variable X has the following probability density function:

$$f(x) = x e^x, \quad 0 < x < 1, \quad \text{zero otherwise.}$$

a) Find $P(X < \frac{1}{2})$.

$$P(X < \frac{1}{2}) = \int_0^{1/2} x e^x dx = \left[x e^x - e^x \right]_0^{1/2} = 1 - \frac{\sqrt{e}}{2} \approx 0.175639.$$

b) Find $\mu = E(X)$.

$$\mu = E(X) = \int_0^1 x \cdot x e^x dx = \left[x^2 e^x - 2x e^x + 2e^x \right]_0^1 = e - 2.$$


c) Find the moment-generating function of X , $M_X(t)$.

$$\begin{aligned} M_X(t) &= \int_0^1 e^{tx} \cdot x e^x dx = \int_0^1 x e^{(t+1)x} dx \\ &= \left[\frac{1}{t+1} x e^{(t+1)x} - \frac{1}{(t+1)^2} e^{(t+1)x} \right]_0^1 \\ &= \frac{1}{t+1} e^{t+1} - \frac{1}{(t+1)^2} e^{t+1} + \frac{1}{(t+1)^2} \\ &= \frac{t e^{t+1} + 1}{(t+1)^2}, \quad t \neq -1. \end{aligned}$$

$$M_X(-1) = \int_0^1 x dx = \frac{1}{2}.$$

11. Let X be a continuous random variable with the probability density function

$$f(x) = \begin{cases} c|x-3|, & 0 < x < 8, \\ 0, & \text{otherwise.} \end{cases}$$



a) Find the value of c that makes $f(x)$ a valid probability density function.

$$\text{Must have } \int_{-\infty}^{\infty} f(x) dx = 1.$$

$$1 = \int_0^8 c|x-3| dx = c \int_0^3 (3-x) dx + c \int_3^8 (x-3) dx = \frac{9}{2}c + \frac{25}{2}c = 17c.$$


$$\Rightarrow 1 = 17c. \quad \Rightarrow c = \frac{1}{17}.$$

b) Find the probability $P(X < 5)$.

$$P(X < 5) = \int_0^5 \frac{1}{17}|x-3| dx = \frac{1}{17} \int_0^3 (3-x) dx + \frac{1}{17} \int_3^5 (x-3) dx = \frac{9}{34} + \frac{4}{34} = \frac{13}{34}.$$

OR

$$P(X < 5) = 1 - P(X \geq 5) = 1 - \int_5^8 \frac{1}{17}|x-3| dx = 1 - \frac{1}{17} \int_5^8 (x-3) dx = 1 - \frac{21}{34} = \frac{13}{34}.$$



c) Find the median of the probability distribution of X .

$$F_X(x) = 0, \quad x < 0,$$

$$F_X(x) = \frac{1}{17} \int_0^x (3-y) dy = \frac{9}{34} - \frac{(3-x)^2}{34}, \quad 0 \leq x < 3,$$

$$F_X(x) = \frac{1}{17} \int_0^3 (3-y) dy + \frac{1}{17} \int_3^x (y-3) dy = \frac{9}{34} + \frac{(x-3)^2}{34}, \quad 3 \leq x < 8,$$


$$F_X(x) = 1, \quad x \geq 8.$$

$$F_X(3) = \frac{9}{34}. \quad \Rightarrow \quad 3 < \text{median} < 8.$$

$$F_X(\text{median}) = \frac{9}{34} + \frac{(\text{median}-3)^2}{34} = \frac{1}{2}.$$

$$\Rightarrow (\text{median}-3)^2 = 8.$$

$$\Rightarrow \text{median} = \mathbf{3 + \sqrt{8}} \approx 5.828427.$$



d) Find the mean of the probability distribution of X.

$$\begin{aligned}
 E(X) &= \int_{-\infty}^{\infty} x \cdot f(x) dx = \frac{1}{17} \int_0^3 x(3-x) dx + \frac{1}{17} \int_3^8 x(x-3) dx = \frac{9}{34} + \frac{475}{102} \\
 &= \frac{502}{102} = \frac{\mathbf{251}}{\mathbf{51}} \approx 4.92157.
 \end{aligned}$$

e) Find the variance of the probability distribution of X.

$$\begin{aligned}
 E(X^2) &= \int_{-\infty}^{\infty} x^2 \cdot f(x) dx = \frac{1}{17} \int_0^3 x^2(3-x) dx + \frac{1}{17} \int_3^8 x^2(x-3) dx = \frac{27}{68} + \frac{2075}{68} \\
 &= \frac{1051}{34} \approx 30.912.
 \end{aligned}$$

$$\text{Var}(X) = E(X^2) - [E(X)]^2 = \frac{1051}{34} - \left(\frac{251}{51}\right)^2 = \frac{\mathbf{34801}}{\mathbf{5202}} \approx 6.69.$$