

p.m.f. or p.d.f. $f(x; \theta)$, $\theta \in \Omega$. Ω – parameter space.

1. Suppose $\Omega = \{ 1, 2, 3 \}$ and the p.d.f. $f(x; \theta)$ is

$$\theta = 1: \quad f(1; 1) = 0.6, \quad f(2; 1) = 0.1, \quad f(3; 1) = 0.1, \quad f(4; 1) = 0.2.$$

$$\theta = 2: \quad f(1; 2) = 0.2, \quad f(2; 2) = 0.3, \quad f(3; 2) = 0.3, \quad f(4; 2) = 0.2.$$

$$\theta = 3: \quad f(1; 3) = 0.3, \quad f(2; 3) = 0.4, \quad f(3; 3) = 0.2, \quad f(4; 3) = 0.1.$$

What is the maximum likelihood estimate of θ (based on only one observation of X) if ...

a) $X = 1;$

b) $X = 2;$

c) $X = 3;$

d) $X = 4.$

Likelihood function:

$$L(\theta) = L(\theta; x_1, x_2, \dots, x_n) = \prod_{i=1}^n f(x_i; \theta) = f(x_1; \theta) \cdot \dots \cdot f(x_n; \theta)$$

It is often easier to consider $\ln L(\theta) = \sum_{i=1}^n \ln f(x_i; \theta).$

Maximum Likelihood Estimator: $\hat{\theta} = \arg \max L(\theta) = \arg \max \ln L(\theta).$

Method of Moments:

$$E(X) = g(\theta). \quad \text{Set } \bar{X} = g(\tilde{\theta}). \quad \text{Solve for } \tilde{\theta}.$$

- 2.** Let X_1, X_2, \dots, X_n be a random sample of size n from a Poisson distribution with mean λ , $\lambda > 0$. That is,

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad k = 0, 1, 2, 3, \dots$$

- a) Obtain the method of moments estimator of λ , $\tilde{\lambda}$.
- b) Obtain the maximum likelihood estimator of λ , $\hat{\lambda}$.

- 3.** Let X_1, X_2, \dots, X_n be a random sample of size n from a Geometric distribution with probability of “success” p , $0 < p < 1$. That is,


$$P(X = k) = (1 - p)^{k-1} p, \quad k = 1, 2, 3, \dots$$

- a) Obtain the method of moments estimator of p , \tilde{p} .
- b) Obtain the maximum likelihood estimator of p , \hat{p} .

4. Let X_1, X_2, \dots, X_n be a random sample of size n from the distribution with probability density function

$$f(x; \theta) = \begin{cases} \frac{1}{\theta} \cdot x^{1-\theta} & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$0 < \theta < \infty.$$



- a) Obtain the method of moments estimator of θ , $\tilde{\theta}$.

Method of Moments:

$$E(X) = g(\theta). \quad \text{Set } \bar{x} = g(\tilde{\theta}). \quad \text{Solve for } \tilde{\theta}.$$

- b) Obtain the maximum likelihood estimator of θ , $\hat{\theta}$.

Likelihood function:

$$L(\theta) = L(\theta; x_1, x_2, \dots, x_n) = \prod_{i=1}^n f(x_i; \theta) = f(x_1; \theta) \cdot \dots \cdot f(x_n; \theta)$$

$$\text{Maximum Likelihood Estimator: } \hat{\theta} = \arg \max L(\theta) = \arg \max \ln L(\theta).$$

4. (continued)

c) Suppose $n = 3$, and $x_1 = 0.2$, $x_2 = 0.3$, $x_3 = 0.5$. Compute the values of the method of moments estimate and the maximum likelihood estimate for θ .

5. Let X_1, X_2, \dots, X_n be a random sample of size n from $N(\theta_1, \theta_2)$, where $\Omega = \{(\theta_1, \theta_2) : -\infty < \theta_1 < \infty, 0 < \theta_2 < \infty\}$. That is, here we let $\theta_1 = \mu$ and $\theta_2 = \sigma^2$.

a) Obtain the maximum likelihood estimator of θ_1 , $\hat{\theta}_1$, and of θ_2 , $\hat{\theta}_2$.

b) Obtain the method of moments estimator of θ_1 , $\tilde{\theta}_1$, and of θ_2 , $\tilde{\theta}_2$.